

リウマチ・膠原病内科グループ 英語論文の業績 2018年

- 1) Tamechika SY, Sasaki K, Hayami Y, Ohmura SI, Maeda S, Iwagaitsu S, Naniwa T. Patient satisfaction and efficacy of switching from weekly bisphosphonates to monthly minodronate for treatment and prevention of glucocorticoid-induced osteoporosis in Japanese patients with systemic rheumatic diseases: a randomized, clinical trial. *Arch Osteoporos*. 13(1):67, 2018.
- 2) Watanabe H, Sada KE, Matsumoto Y, Harigai M, Amano K, Dobashi H, Fujimoto S, Usui J, Yamagata K, Atsumi T, Banno S, Sugihara T, Arimura Y, Matsuo S, Makino H; Japan Research Committee of the Ministry of Health, Labour, and Welfare for Intractable Vasculitis, and the Research Committee of Intractable Renal Disease of the Ministry of Health, Labour, and Welfare of Japan. Association Between Reappearance of Myeloperoxidase-Antineutrophil Cytoplasmic Antibody and Relapse in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Subgroup Analysis of Nationwide Prospective Cohort Studies. *Arthritis Rheumatol*. 70(10):1626-1633, 2018.
- 3) Sato S, Masui K, Nishina N, Kawaguchi Y, Kawakami A, Tamura M, Ikeda K, Nunokawa T, Tanino Y, Asakawa K, Kaneko Y, Gono T, Ukichi T, Kaieda S, Naniwa T, Kuwana M; JAMI investigators . Initial predictors of poor survival in myositis-associated interstitial lung disease: a multicentre cohort of 497 patients. *Rheumatology (Oxford)*. 2018.
- 4) Naniwa T, Iwagaitsu S, Kajiura M. Long-term efficacy and safety of add-on tacrolimus for persistent, active rheumatoid arthritis despite treatment with methotrexate and tumor necrosis factor inhibitors. *Int J Rheum Dis*. 21(3):673-687, 2018.